



May-August 2013 | Vol.32 No.2

UNIVERSA MEDICINA

<http://www.univmed.org>

- *Phaleria macrocarpa* reduces glomerular growth factor expression in alloxan-induced diabetic rats
 - Antifungal activity of neem leaf ethanol extract on *Aspergillus flavus*
- Red fruit oil supplementation fails to prevent oxidative stress in rats
 - Nicotine supplementation blocks oocyte maturation in *Rattus norvegicus*
 - *Andrographis paniculata* extract induced apoptosis of adenocarcinoma mammae in C3H mice
- Weekly lifestyle counselling improves glucose level in type 2 diabetes mellitus patients
- Soy-isoflavone supplementation tends to reduce menopausal symptoms in postmenopausal women
- *Plasmodium falciparum* infection and the risk of anemia in school children

ISSN 1907-3062

UNIVERSA MEDICINA



Editor-in-Chief

Prof Adi Hidayat, MD, MS, PhD

Manuscripts Editor

Prof Murad Lesmana, MD

Editorial Board

Prof H.A.Prayitno, MD, SpKJ, PhD

Prof Muzief Munir, MD, SpA(K)

Prof Julius E. Surjawidjaja, MD, SpMK

Raditya Wratsangka, MD, SpOG(K), PhD

Oktavianus CH. Salim, MD, MS

Elly Herwana, MD, MBIomed, PhD

Pusparini, MD, SpPK, PhD

Maria Regina Rachmawati, MD, SpRM, PhD

Meiyanti, MD, SpFK

Yenny, MD, SpFK

Richard Tjan, MD, DTMH

Distribution and Marketing

Eddy Kasim, DS

Julius I. Mulia, MD

Secretary

Rita Hemawati

Accreditation : 58/DIKTI/Kep/2013

Correspondence Address:

Medical Faculty, Trisakti University

Jl. Kyai Tapa No.260

Grogol - Jakarta 11440

Phone : 021-5672731 ext. 2611

Fax : 021-5660706

Homepage : www.univmed.org

Email : editors@univmed.org

Published by Faculty of Medicine Trisakti University

Guidelines For Authors

Universa Medicina is a quarterly medical journals and accepts medical and health-related original articles (research papers) in English or Indonesia. The Indonesian papers will be translated and published in English

SUBMISSION

Submitted manuscripts should not have been published before and must not be under simultaneous consideration by any other journal.

Manuscript preparation

The manuscript should be formatted as follows: A4 (212 x 297 mm), with margins of at least 2.5 cm; use double-spacing in a serif font (e.g. Times), 12-point and limited to approximately 16 pages in length including references, tables and figures. Do not justify the right margin. Number pages consecutively in the upper right-hand corner of each page, beginning with the title page. Each manuscript component should begin on a new page in the following sequence: title page, abstract and key words, text, acknowledgments, references, tables; each table on a separate page, complete with title and footnotes, legends for figures. All manuscripts should be accompanied by a cover letter from the author responsible for correspondence.

Title page

This should carry the title of the article, the names and addresses of all authors (the institution to which the work is to be attributed should be listed first), the name, address, fax number and email address of the corresponding author, and a short running title for page headings not more than 40 characters.

Abstract and key words

Original research articles must include a structured abstract that contains no more than 250 words, and should consist of background including objective, methods, design, results and conclusions. Below the abstract, provide a list of 3–10 key words.

Text

The text of research papers should be divided into sections with the following headings: Introduction, Methods, Results, Discussion and Conclusions.

Acknowledgements

Acknowledge only persons who have made substantive contributions to the study.

References

It is the authors' responsibility to check all references very carefully for accuracy and completeness. Number references consecutively in the order in which they are first mentioned in the text. Identify references in the text, Tables and Legends by Arabic numerals in superscript and parenthesis. "Unpublished observations" and "personal communications" may not be used as references; if cited, a letter (from the person quoted) granting permission must be submitted. Authors

should avoid using abstracts as references. Abbreviate journal names according to the Index Medicus system (Also see International Committee of Medical Journal Editors Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Sample References http://www.nlm.nih.gov/bsd/uniform_requirements.html). Examples of correct references are given at the end of these instructions.

Tables and Illustrations (Figures)

Type or print each table with double spacing on a separate sheet of paper. Number tables consecutively in the order of their first citation in the text and supply a brief title for each. Do not use internal horizontal or vertical lines. Give each column a short or abbreviated heading. Authors should place explanatory matter in footnotes, not in the heading. Explain in footnotes all nonstandard abbreviations. For footnotes use the following symbols, in sequence: *, †, ‡, §, ||, ¶, **, ††, ‡‡. Identify statistical measures of variations, such as standard deviation and standard error of the mean. Figures should be either professionally drawn and photographed, or submitted as photographic quality digital prints. For x-ray films, scans, and other diagnostic images, as well as pictures of pathology specimens or photomicrographs, send sharp, glossy, black-and-white or color photographic prints, usually 127 x 173 mm (5 x 7 inches). Figures should be numbered consecutively according to the order in which they have been first cited in the text. If a figure has been published, acknowledge the original source and submit written permission from the copyright holder to reproduce the material. Permission is required irrespective of authorship or publisher except for documents in the public domain.

Units of Measurement

Measurements of length, height, weight, and volume should be reported in metric units (meter, kilogram, or liter) or their decimal multiples. Temperatures should be in degrees Celsius. Blood pressures should be in millimeters of mercury, unless other units are specifically required by the journal.

Sending the Manuscript to the Journal

Manuscripts, written in English, can be submitted electronically as an e-mail attachment to the following website: <http://www.univmed.org> or alternatively to the following address:

Editor's Office, Universa Medicina
Fakultas Kedokteran Universitas Trisakti
Kampus B USAKTI Lt. 6
Jl. Kyai Tapa No. 260
Grogol-Jakarta Barat 11440
Phone (62-021) 5672731 (hunting) ext 2504 Fax 62-021-5660706
Email: editors@univmed.org

Examples of correct forms of references

Journals

1. Standard journal

List all authors when 6 or fewer; when 7 or more, list only the first 6 and add "et al." Abbreviate journal titles according to Index Medicus style, which is used in MEDLINE citations.

Pinelli NR, Hurren KM. Efficacy and safety of long-acting glucagon-like peptide-1 receptor: a systematic review and meta-analysis. *Ann Pharmacother* 2011;45:850-60.

Campbell NRC, Gilbert RE, Leiter LA, Larochelle P, Tobe S, Chokalingam A, et al. Hypertension in people with type 2 diabetes: update on pharmacologic management. *Can Fam Physician* 2011;57:997-1002.

2. Corporate author

Diabetes Prevention Program Research Group. Hypertension, insulin, and proinsulin in participants with impaired glucose tolerance. *Hypertension* 2002;40:679-86.

3. Volume with supplement

Geraud G, Spierings EL, Keywood C. Tolerability and safety of frovatriptan with short- and long-term use for treatment of migraine and in comparison with sumatriptan. *Headache* 2002;42 Suppl 2:S93-9.

4. Electronic journal without page numbers

Santos CAST, Fiaccone RL, Oliveira NF, Cunha S, Barreto ML, do Carmo BB, et al. Estimating adjusted prevalence ratio in clustered cross-sectional epidemiological data. *BMC Med Res Method* 2008;8:80 doi:10.1186/1471-2288-8-80.

Thyssen JP, Linneberg A, Carlsen BC, Johansen JD, Engkilde K, Hansen T, et al. A possible association between a dysfunctional skin barrier (filaggrin null-mutation status) and diabetes: a cross-sectional study. *BMJ Open* 2011 doi:10.1136/bmjopen-2011-000062.

Bawaskar HS, Bawaskar PH. Efficacy and safety of scorpion antivenom plus prazosin compared with prazosin alone for venomous scorpion (*Mesobuthus tamulus*) sting: randomised open label clinical trial. *BMJ* 2011;342:c7136. doi:10.1136/bmj.c7136.

Books and Other Monographs

1. Editor(s), compiler(s) as author

Gilstrap LC, Cunningham FG, VanDorsten JP, editors. *Operative obstetrics*. 2nd ed. New York: McGraw-Hill;2002.

2. Chapter in a book

Meltzer PS, Kallioniemi A, Trent JM. Chromosome alterations in human solid tumors. In: Vogelstein B, Kinzler KW, editors. *The genetic basis of human cancer*. New York: McGraw-Hill;2002.p.93-113.

3. Conference paper

Christensen S, Oppacher F. An analysis of Koza's computational effort statistic for genetic programming. In: Foster JA, Lutton E, Miller J, Ryan C, Tettamanzi AG, editors. *Genetic programming. EuroGP 2002: Proceedings of the 5th European Conference on Genetic Programming*; 2002 Apr 3-5; Kinsdale, Ireland. Berlin: Springer;2002.p.182-91.

4. Dissertation

Borkowski MM. Infant sleep and feeding: a telephone survey of Hispanic Americans [dissertation]. Mount Pleasant (MI): Central Michigan University;2002.

Electronic Material

1. CD-ROM

Anderson SC, Poulsen KB. Anderson's electronic atlas of hematology [CD-ROM]. Philadelphia: Lippincott Williams & Wilkins;2002.

2. Journal article on the internet

Rosenthal S, Chen R, Hadler S. The safety of a cellular pertussis vaccine vs whole cell pertussis vaccine. Available at: http://www.ama-assn.org/sci_publ/journals/archive/ajdc/vol_150/no_5/full/htm. Accessed Jun 10, 2007.

3. Monograph on the internet

Foley KM, Gelband H, editors. *Improving palliative care for cancer* [monograph on the Internet]. Washington: National Academy Press; 2001. Available at: <http://www.nap.edu/books/0309074029/html/>. Accessed July 9, 2002.

Manuscript Checklist

As you prepare to submit your manuscript, please go through this list and check-off each as it is completed. Failure to conform to the format requirements will result in your paper being sent back to you, and a delay in publication.

Include original manuscript plus two copies and a CD containing an exact version of the original hard copy

Double space manuscripts (text, references and tables) and leave right margins unjustified (ragged)
The title is in upper case letters and centered on the page. The author's names are centered under the title

Information about the author's department, university, university address, and e-mail address is provided. Phone and fax numbers appear only on the cover letter. If information is different for each author, multiple asterisks are used

Provide a structured abstract that conforms with the required abstract format

An acknowledgement section can be presented after the conclusion

References are numbered and listed in the reference section in the order that they appear in the text, not in alphabetical order by author's names. When referring to a source in the text, only the numeral enclosed in square brackets is used for identification.

Pedoman Untuk Penulis

Universa Medicina adalah berkala ilmiah yang terbit setiap catur wulan (3 kali dalam 1 tahun, Januari-April, Mei-Agustus, September-Desember) dan menerima tulisan ilmiah di bidang kesehatan/kedokteran dalam bentuk laporan penelitian (*original article/research paper*) dalam bahasa Indonesia atau Inggris. Setiap makalah akan dipublikasi dalam bahasa Inggris.

Pengiriman Makalah

Makalah yang dikirim untuk dimuat dalam Universa Medicina belum pernah dipublikasi dan tidak dikirimkan ke penerbitan lain pada waktu yang bersamaan. Naskah dikirimkan dalam bentuk *print out* sebanyak rangkap 2 (dua), dalam *file* bentuk CD-R/CD-RW, serta diketik menggunakan program komputer *Microsoft Word for Windows* baik melalui e-mail (editor@univmed.org) maupun ke alamat redaksi.

Persiapan teknis makalah

Naskah diketik pada kertas berukuran 8,27 x 11,69" (A4), dengan batas tepi (*margin*) 1", *font* Times New Roman, besar huruf (*font size*) 12 *point* dan menggunakan spasi rangkap 2 (dua) (*double space*). Setiap bagian/komponen dari naskah dimulai pada halaman baru, dengan urutan sebagai berikut: halaman judul, abstrak dan kata kunci (*key words*), teks keseluruhan, ucapan terima kasih, daftar pustaka, tabel dan gambar (setiap tabel dan gambar pada halaman terpisah). Nomor halaman dicantumkan secara berurutan dimulai dari halaman judul pada sudut sebelah kanan bawah. Makalah hendaknya ditulis maksimal 16 halaman.

Halaman judul

Halaman judul (halaman pertama) harus mencakup a) judul makalah yang dibuat sesingkat mungkin, spesifik informatif dan ringkasan judul (*a short running head or foot line*) tidak lebih dari 40 karakter (hitung huruf dan spasi) yang dicantumkan di bawah judul; b) nama dan alamat setiap penulis, nama departemen dan lembaga afiliasi penulis; c) nama dan alamat penulis untuk korespondensi serta nomor telepon, nomor faksimili, dan alamat e-mail.

Abstrak dan kata kunci

Halaman kedua memuat abstrak yang tidak lebih dari 250 kata untuk laporan penelitian yang ditulis dalam bahasa Indonesia dan Inggris. Abstrak laporan penelitian merupakan abstrak terstruktur yang berisikan latar belakang termasuk tujuan penelitian (*background*), metode (*methods*), hasil (*results*) dan kesimpulan (*conclusions*). Kata kunci dicantumkan di bawah abstrak pada halaman yang sama sebanyak 3-10 kata. Gunakanlah kata-kata yang sesuai dengan daftar pada *Index Medicus*.

Teks

Teks makalah laporan penelitian dibagi dalam beberapa bagian dengan judul sebagai berikut: Pendahuluan (*Introduction*), Metode (*Methods*), Hasil (*Result*), Pembahasan (*Discussion*), Kesimpulan (*Conclusion*), dan Ucapan terima kasih (*Acknowledgement*).

Ucapan terima kasih

Terutama ditujukan kepada 1) pihak-pihak yang memberikan bantuan dana dan dukungan, 2) dukungan dari bagian dan lembaga, 3) para profesional yang memberikan kontribusi dalam penyusunan makalah.

Daftar Pustaka

Daftar pustaka ditulis sesuai dengan cara penulisan menurut Vancouver dan hanya mencantumkan kepastaknaan yang dipakai dan relevan. Rujukan diberi nomor urut sesuai dengan penampilannya dalam teks dengan menggunakan angka Arab dan dalam teks nomor urut dituliskan dalam tanda kurung. Acuan yang digunakan hendaknya > 80% merupakan sumber acuan primer (jurnal) dan mutakhir (10 tahun terakhir). Tabel dan gambar diberi nomor sesuai dengan urutan penampilannya dalam teks menggunakan angka Arab. Singkatan nama jurnal ditulis sesuai dengan daftar dalam *Index Medicus* yang dapat diperoleh pada *library's web site* (<http://www.nlm.nih.gov>). Hindari penggunaan abstrak sebagai rujukan.

Contoh cara menuliskan rujukan adalah sebagai berikut :

Artikel dalam jurnal

1. Artikel jurnal standar
Cantumkan semua penulis bila jumlah penulis 6 (enam) atau kurang, bila 7 (tujuh) atau lebih cantumkan 6 (enam) penulis pertama dan diikuti oleh et al.
Pinelli NR, Hurren KM. Efficacy and safety of long-acting glucagon-like peptide-1 receptor: a systematic review and meta-analysis. *Ann Pharmacother* 2011;45:850-60.
Lebih dari 6 (enam) penulis :
Campbell NRC, Gilbert RE, Leiter LA, Larochelle P, Tobe S, Chokalingam A, et al. Hypertension in people with type 2 diabetes: update on pharmacologic management. *Can Fam Physician* 2011;57: 997-1002.
2. Artikel jurnal yang disertai DOI
Bhutta ZA, Darmstadt GL, Hasan BS, Haws RA. Community-based interventions for improving perinatal and neonatal health outcomes in developing countries: a review of the evidence. *Pediatrics* 2005;115 Suppl 2:519-617. DOI:10.1542/peds.2004-1441.
3. Organisasi sebagai penulis
The National Osteoporosis Foundation of South Africa. Use of generic alendronate in the treatment of osteoporosis. *S Afr Med J* 2006;96:696-7.
4. Volume dengan suplemen
Geraud G, Spierings EL, Keywood C. Tolerability and safety of frovatriptan with short- and long-term use for treatment of migraine and in comparison with sumatriptan. *Headache* 2002;42 Suppl 2:S93-9.
5. Edisi dengan suplemen
Glauser TA. Integrating clinical trial data into clinical practice. *Neurology* 2002;58(12 Suppl 7):S6-12.
6. Volume dengan bagian
Abend SM, Kulish N. The psychoanalytic method from an epistemological viewpoint. *Int J Psychoanal* 2002;83(Pt 2):491-5.
7. Edisi dengan bagian
Ahrar K, Madoff DC, Gupta S, Wallace MJ, Price RE, Wright KC. Development of a large animal model for lung tumors. *J Vasc Interv Radiol* 2002;13(9 Pt 1):923-8.
8. Edisi tanpa volume
Banit DM, Kaufer H, Hartford JM. Intraoperative frozen section analysis in revision total joint arthroplasty. *Clin Orthop* 2002;401:230-8.

9. Nomor Halaman dalam angka romawi
Chadwick R, Schuklenk U. The politics of ethical consensus finding. *Bioethics* 2002;16:iii-v.

Buku dan monograf lain

(Aturan Vancouver sebelumnya salah mencantumkan tanda baca koma dan bukannya tanda baca titik koma antara penerbit dan tahun penerbitan)

1. Penulis perorangan
Murray PR, Rosenthal KS, Kobayashi GS, Pfaffler MA. *Medical microbiology*. 4th ed. St. Louis: Mosby;2002.
2. Editor, sebagai penulis
Jacobson A, Jacobson RL, editors. *Radiographic cephalometry from basic to 3-D imaging*, 2nd ed. New Maiden: Quintessence;2006.
3. Organisasi sebagai penulis dan penerbit
Ministry of Health, Republic of Indonesia. *Behavioral surveillance survey (BSS) result in Indonesia 2004-2005*. Jakarta: Ministry of Health;2005.
4. Meltzer PS, Kallioniemi A, Trent JM. Chromosome alterations in human solid tumors. In: Vogelstein B, Kinzler KW, editors. *The genetic basis of human cancer*. New York: McGraw-Hill;2002.p.93-113.
5. Bab dalam buku
Catatan : Aturan Vancouver sebelumnya mencantumkan tanda baca titik dua, sekarang tanpa 'p' untuk halaman.
Swierkosz EM, Arens MQ. Susceptibility test methods: viruses. In: Murray PR, Baron EJ, Jorgensen JH, Pfaffler MA, Tenover FC, Tenover RH, editors. *Manual of clinical microbiology*, vol. 2. 8th ed. Washington DC: American Society for Microbiology;2003.p.706-18.
6. Prosiding konferensi
Dostrovsky JO, Carr DB, Koltzenburg M, editors. *Proceedings of the 10th World Congress on Pain*; 2002 Aug 17-22; San Diego, CA. Seattle:IASP Press;2003.
7. Makalah dalam konferensi
Antani S, Long LR, Thoma GR, Lee DJ. Anatomical shape representation in spine x-ray images. *Proceedings of the 3rd IASTED International Conference on Visualization, Imaging and Image Processing*; Sep 8-10; Benalmadena, Spain;2003.
8. Laporan ilmiah atau teknis
Diterbitkan oleh agen penyandang dana/sponsor:
Yen GG (Oklahoma State University, School of Electrical and Computer Engineering, Stillwater, OK). *Health monitoring on vibration signatures. Final report*. Arlington (VA): Air Force Office of Scientific Research (US), Air Force Research Laboratory; 2002. Rweport No: D AFRISRLTR020123. Contract No.: F496209810049
Diterbitkan oleh agen pelaksana:
Russell ML, Goth-Goldstein R, Apte MG, Fisk WJ. *Method for measuring the size distribution of airborne Rhinovirus*.

Berkeley (CA): Lawrence Berkeley National Library, Environmental Energy Technologies Division;2002. Report No.: LBNL49574. Contract No.:DEAC0376SF00098. Sponsored by Department of Energy.

9. Disertasi
Borkowski MM. *Infant sleep and feeling: a telephone survey of Hispanic Americans [dissertation]*. Mount Pleasant (MI): Central Michigan University;2002.
10. Materi audiovisual
Chason KW, Sallustio S. *Hospital preparedness for bioterrorism [videocassette]*. Secaucus (NJ): Network for Continuing Medical Education;2002.
11. Kamus, Encyclopedia, dan rujukan serupa
Harahan C, Valerian. In: Krapp K, Longe JL, editors. *The gate encyclopedia of alternative medicine*. Michigan: Gate Group;2001. vol 4, p.17668-70.

Materi Elektronik

1. Artikel jurnal dalam format elektronik
Barr AE, Barbe MF. Pathophysiological tissue changes associated with repetitive movement : a review of the evidence. *Phys Ther* 2002;82:173-87. Available at: <http://http://www.physiotherapy.org/content/82/2/173.full.pdf+html>. Accessed August 10, 2011.
2. Monograf dalam format elektronik
Anderson SC, Poulsen KB. *Anderson's electronic atlas of hematology [CD-ROM]*. Philadelphia: Lippincott Williams & Wilkins; 2002.

Tabel dan gambar

Cetak setiap tabel pada halaman terpisah dan diketik spasi rangkap dua (*double space*). Nomor urut tabel dan gambar sesuai dengan urutan penampilannya dalam teks. Untuk catatan kaki (*footnotes*) pada tabel gunakan symbol dengan urutan sebagai berikut : *, †, ‡, §, ||, ¶, **, ††, ‡‡.

Naskah yang diterima redaksi akan dibahas oleh pengasuh dan redaksi berhak memperbaiki susunan bahasa tanpa mengubah isinya. Penggunaan istilah asing non-medis sedapat mungkin dihindari atau disertai terjemahan penjelasannya. Usulan perbaikan naskah (terutama menyangkut substansi) akan disampaikan kepada penulis yang bersangkutan.

Naskah dikirimkan ke alamat Redaksi :
Fakultas Kedokteran Universitas Trisakti
Jl. Kyai Tapa No. 260, Grogol - Jakarta 11440
Telepon: (021) 5672731 (hunting) ext. 2054. Fax. (021) 5660706
E-mail: editors@univmed.org

Tiras : 800 eks

Table of Contents

Volume 32 — May-August, 2013 — Number 2

	<u>Pages</u>
Editorial	
HIV antiretroviral preexposure prophylaxis..... <i>Richard Tjan</i>	69
Research Articles	
<i>Phaleria macrocarpa</i> reduces glomerular growth factor expression in alloxan-induced diabetic rats..... <i>Evy Sulistyoningrum and Setiawati</i>	71
Antifungal activity of neem leaf ethanol extract on <i>Aspergillus flavus</i> <i>Ayly Margaret, Hanna Yolanda, and Lies K. Wibisono</i>	80
Red fruit oil supplementation fails to prevent oxidative stress in rats..... <i>Maria Dara Novi Handayani, Parwati Abadi Soekarno, and Septelia Inawati Wanandi</i>	86
Nicotine supplementation blocks oocyte maturation in <i>Rattus norvegicus</i> <i>Meitria Syahadatina Noor, H.M. Bakhriansyah, Widjiati, and Budi Santoso</i>	92
<i>Andrographis paniculata</i> extract induced apoptosis of adenocarcinoma mammae in C3H mice..... <i>Nugrahaningsih, Sarjadi, Edi Dharmana, and Hertanto Wahyu Subagio</i>	99
Weekly lifestyle counselling improves glucose level in type 2 diabetes mellitus patients <i>Amallia Nuggetsiana Setyawati, Inggar Octa Pusthika, and Kusmiyati Tjahjono DK</i>	108
Soy-isoflavone supplementation tends to reduce menopausal symptoms in postmenopausal women <i>Raditya Wratsangka and Atut Cicih Mayasari</i>	118
<i>Plasmodium falciparum</i> infection and the risk of anemia in school children..... <i>Suriyani Tan, Taniawati Supali, and Heri Wibowo</i>	128

Editorial

HIV antiretroviral preexposure prophylaxis

Richard Tjan
Editor

According to 3 field trials conducted in Africa, one among African women and two among heterosexual couples, antiretroviral preexposure prophylaxis for prevention of HIV-1 has been shown to be effective.⁽¹⁻³⁾ In preexposure prophylaxis, persons without HIV infection are given an oral drug before they have sexual contact with HIV-infected partners.⁽⁴⁾ The drug in question is tenofovir disoproxil fumarate (TDF), a prodrug of tenofovir.⁽¹⁾ On the basis of the 3 field trials, the Antiviral Drugs Advisory Committee of the Food and Drug Administration has recommended a combination of antiretroviral drugs (tenofovir/emtricitabine) for preexposure prophylaxis of HIV.⁽⁵⁾

TDF is currently indicated for the treatment of HIV in adults over 18 years of age or hepatitis B virus (HBV) infection in adults, or both. The drug is called a nucleotide reverse transcriptase inhibitor (NRTI), preventing the synthesis of viral copies by HIV reverse transcriptase or HBV DNA polymerase.⁽⁶⁾ To retard the emergence of TDF resistance, the drug is usually given in combination with another antiviral, such as emtricitabine (FTC). Nucleic acid testing for HIV virus when starting preexposure prophylaxis, may reduce the risk of resistance, but it is at present not an option in developing countries because of its high costs. Rare but potentially serious adverse reactions to TDF are lactic acidosis and toxic effects on the liver and kidneys.⁽⁶⁾ Because administration of TDF to healthy noninfected persons implies using the drug for a prolonged period of many years, the long-term safety of TDF and the TDF-FTC combination has to be clearly established.⁽⁴⁾

There is also a real possibility that preexposure prophylaxis may lead to relaxation of the customary precautions on the part of the sexual partners, such as engaging in increased risky sexual behavior or abandoning the use of conventional prophylactic measures (e.g. condoms).⁽⁴⁾ This matter should be a problem for health educators.

From a practical point of view, because of the potential of serious liver and kidney disease caused by TDF, the medical practitioner should prescribe preexposure prophylaxis only in high risk cases, and not for prevention of HIV in otherwise healthy individuals, e.g. blood bank personnel or dental practitioners with a low risk of exposure to HIV, which are currently not indicated. Prescription should be done on an individual basis.

Indeed, the old Hippocratic advice of not too readily prescribing any new modes of treatment, or in plain words - Wait and watch- still holds true. This is presumably one of the reasons for not blindly or overenthusiastically accepting HIV preexposure prophylaxis.

REFERENCES

1. Van Damme L, Corneli A, Ahmed K, Agot K, Lombaard J, Kapiga S, et al. for the FEM-PrEP Study Group. Preexposure prophylaxis for HIV infection among African women. *N Engl J Med* 2012;367:411-22.
2. Thigpen MC, Kebaabetswe PM, Paxton LA, Smith DK, Rose CE, Segolodi TM, et al. for the TDF2 Study Group. Antiretroviral preexposure prophylaxis for heterosexual HIV transmission in Botswana. *N Engl J Med* 2012;367:423-34.
3. Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, Wangisi J, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *N Engl J Med* 2012;367:399-410.
4. Cohen MS, Baden LR. Preexposure prophylaxis for HIV - where do we go from here? *N Engl J Med* 2012;367:459-61.
5. Mascolini M. FDA panel endorses Truvada for pre-exposure prophylaxis. International AIDS Society, May 12, 2012. Available at: <http://www.iasociety.org/Article.aspx?elementId=14505>. Accessed 13 August 2013.
6. World Health Organization. Patient information leaflet. WHOPAR part 3 11/2010, version 1.0.

Nicotine supplementation blocks oocyte maturation in *Rattus norvegicus*

Meitria Syahadatina Noor*, H.M. Bakhriansyah**, Widjiati***,

*Department of Public Health,
Faculty of Medicine,
University of Lambung
Mangkurat, Banjarmasin

**Department of Pharmacology,
Faculty of Medicine,
University of Lambung
Mangkurat, Banjarmasin

***Department of Embryology,
Faculty of Veterinary
Medicine, University of
Airlangga, Surabaya

****Department of Obstetrics and
Gynecology, Faculty of
Medicine, Dr. Soetomo General
Hospital, University of
Airlangga, Surabaya

Correspondence dr. Meitria
Syahadatina Noor Department
of Public Health,
Faculty of Medicine,
University of Lambung
Mangkurat
Jl. Veteran, Banjarmasin,
Kalimantan Selatan
Phone: +6281391739795
Email: drmeitria79@gmail.com

Univ Med 2013;32:92-8

**and Budi
Santoso******

after China and India. Nicotine as the main component of cigarette smoke has negative effects on the reproductive system, such as oocyte maturation, ovulation, and fertilization, and increasing the diploidy of oocytes. The goal of this research was to evaluate the effect of nicotine on oocyte maturation in *Rattus norvegicus*.

METHODS

This was an experimental study with post test only control group design. The subjects were 40 rats selected homogenously and randomly. They were divided into a control group (receiving carboxy-methyl-cellulose sodium and 3 treatment groups (I-III) receiving nicotine subcutaneously for 7 days at dosages of 21 mg/ kgBW, 41 kg/kgBW and 84/kgBW, respectively. The observations comprised oocyte maturation stage, viz. germinal vesicle (GV), germinal vesicle breakdown (GVBD), metaphase I and metaphase II. Data were analyzed by one-way Anova with $\alpha=0.05$, followed by Tukey's HSD test.

RESULTS

One-way Anova showed significant differences in oocyte maturation in all groups. Tukey's HSD test showed that for GV, the differing groups were control and I, control and II, I and III. For GVBD, the differing groups were control and I, I and II, I and III. For metaphase I, the differing groups were control with I, II, and III, I and II, I and III. For metaphase II, the differing groups were control versus I, II, and III, I and II, I and III.

CONCLUSION

Low dose of nicotine is capable of affecting oocyte maturation in *Rattus norvegicus*.

Key words: Nicotine, maturation, oocyte, *Rattus norvegicus*

ABSTRACT

BACKGROUND

Indonesia has the third
largest tobacco
consumption in the world

Pemberian nikotin menghambat maturasi oosit pada Rattus norvegicus

ABSTRAK

LATAR BELAKANG

Indonesia adalah negara konsumen tembakau terbesar di dunia dan menempati urutan ketiga setelah Cina dan India. Nikotin sebagai komponen terbesar dari rokok memiliki efek terhadap sistem reproduksi, antara lain dapat menghambat maturasi oosit, menurunkan ovulasi dan fertilisasi, serta meningkatkan jumlah oosit yang diploidi. Tujuan penelitian ini adalah untuk menilai efek nikotin terhadap maturasi oosit pada *Rattus norvegicus*.

METODE

Penelitian ini adalah penelitian eksperimental dengan post test only control group design dengan menggunakan 40 ekor tikus putih. Kelompok hewan coba secara acak dibagi menjadi kontrol diberikan carboxy-methyl-cellulose sodium, kelompok I-III yang masing-masing diberikan injeksi nikotin subkutan selama 7 hari dengan dosis 21 mg/kgBB, 42 mg/kgBB, dan 84 mg/kgBB. Hasil yang diamati adalah germinal vesicle (GV), germinal vesicle breakdown (GVBD), metafase I dan II. Anova satu arah dengan taraf kepercayaan 95% digunakan untuk membedakan anatar keempat kelompok, bila berbeda bermakna maka dilanjutkan dengan uji Tukey HSD.

HASIL

Ada perbedaan semua fase maturasi oosit antara keempat kelompok ($p < 0,05$). Hasil uji Tuckey HSD untuk fase GV menunjukkan terdapat perbedaan antara kontrol dan I, kontrol dan II, I dan II, I dan III, serta II dan III. Pada fase GVBD, terdapat perbedaan antara kontrol dan I, I dan II, serta I dan III. Pada fase metaphase I, terdapat perbedaan antara kontrol dengan I, II, dan III, I dan II, serta I dan III. Sedangkan pada fase metaphase II terdapat perbedaan antara kontrol dengan I, II, dan III, I dan II, serta I dan III.

KESIMPULAN

Pemberian nikotin dalam dosis rendah mampu menghambat maturasi oosit pada *Rattus norvegicus*.

Kata kunci: Nikotin,
maturasi, oosit, tikus
putih

around one billion people died from smoking in the 21st century.⁽¹⁾ Nicotine is the main component of cigarette smoke and the cause of the smoking habit or addiction.⁽²⁾

Exposure to cigarette smoke affects both active and passive smokers.⁽³⁾ The effects of cigarette smoke on the reproductive system are to influence the production and function of gametes, ovulation, the reproductive cycle, fertilization, and embryo transport and implantation.⁽⁴⁾ The main component of cigarettes is nicotine ($C_{10}H_{14}N_2$), which constitutes 50% of all components.⁽⁵⁾ Nicotine is capable of forming free radicals, thus being a pro-oxidant. Nicotine exerts adverse effects on follicle growth, number of follicles, thickness of the endometrium and uterine glands.⁽⁶⁾ It also adversely affects cumulus cells and the organization of microtubules and microfilaments in the oocyte during meiosis. In vitro experiments showed that nicotine induced meiotic blockage in metaphase-I of mouse oocytes, while administration of nicotine to mice in vivo resulted in reduced numbers of ovulated oocytes.⁽⁷⁾

INTRODUCTION

According to a statement by the WHO there were 1.3 billion smokers in the world in 2003 and their numbers will increase to 1.7 billion in 2010. It has been estimated that

The study by Dwirahayu found that nicotine blocks oocyte maturation in *Rattus norvegicus* at dosages of 35, 52.5, and 70 mg/kgBW.

This is because nicotine reduces the size of ovarian follicles, thus affecting the oocyte maturation process.⁽⁸⁾

In view of the above, it was thought necessary to conduct further studies on the effects of nicotine on oocyte maturation, using lower doses of nicotine, in order to determine the minimal dose capable of affecting fertility in females. The doses used were based on those of Kakisina who used doses of 3, 6, and 12 mg/ kgBW for determining developmental abnormalities in mouse embryos. The doses had been validated in preliminary studies and were far below the lethal dose.⁽⁹⁾ These doses were then converted into doses of 21, 42, and 84 mg/ kgBW for use in the Norway rat (*Rattus norvegicus*).

METHODS

Design of the study

This study used an experimental laboratory method with post test only control group design. It was conducted at the Univ Med 32 No.2

Embryology Laboratory, Faculty of Veterinary Medicine, University of Airlangga, Surabaya, from June to July 2011.

Experimental animals

The study subjects were adult female Norway rats (*Rattus norvegicus*). The rats were randomly assigned to one control group and three treatment groups (I-III). The sample size was 9 rats per group, based on $\alpha=0.05$, $\beta=0.2$ and effect size = 0.3.⁽¹¹⁾ To anticipate a reduction in numbers through death, the size of the groups was increased by 20%, so that each group contained 11 rats.

Preparation of nicotine

The nicotine doses of 21 mg/kgBW, 42 mg/ kgBW and 84 mg/kgBW, were adjusted to the weight of individual rats. Liquid nicotine of 70% purity was diluted with twice-distilled water (*aqua bidestillata*) and the calculated dose for each rat was administered.

Treatment

The rats were given a subcutaneous injection of nicotine at 21 mg/kgBW (I), 42 mg/kgBW (II) and 84 mg/kgBW (III) for 7 days. These doses were seven times larger than the corresponding dose for mice (7 x mouse dose).⁽¹⁰⁾ The controls were given an injection of carboxy-methyl-cellulose sodium (CMC-Na) using a comparable technique and duration of treatment as used for the treatment groups. The injections were performed by experienced personnel using disposable syringes and needles for each injection.

Harvesting of oocytes

The rats were given an injection of 10 IU of pregnant mare serum gonadotropin (PMSG), and 48 hours later an injection of 10 IU human chorionic gonadotropin (HCG). Subsequently the rats were mated with vasectomized male rats to induce ovulation. After 17 hours each female was examined for the presence of a vaginal plug. Rats with a vaginal plug were sacrificed by euthanasia for harvesting of oocytes.

Harvesting of oocytes was done by lifting the uterus under a dissecting microscope and looking for the fertilization pouch. The oocytes were released by rupturing the fertilization pouch. The released oocytes were then transferred by means of a modified pipette into a petri dish containing phosphate buffered saline (PBS) as washing medium.

Staining of oocytes

The oocytes were placed on a glass slide that was ringed with vaseline and covered with a cover slip. Subsequently the oocytes were fixed in fixing solution (acid alcohol : absolute methanol = 1:3) for at least 24 hours. The slide was then removed from the fixing solution and air-dried. The oocytes were stained in 1% aceto-orcein for 2-3 seconds and washed in decolorizing solution. They were examined under the microscope for germinal vesicles (GV), germinal vesicle breakdown (GVBD), metaphase I and metaphase II. Examination of

Vol.

the oocytes was done at the Embryology Laboratory, Faculty of Veterinary Medicine, University of Airlangga, Surabaya.

Medicine, University of Lambung Mangkurat, Banjarmasin.

RESULTS

The total number of oocytes collected in each treatment group are presented in Figure 1.

Data analysis

The data on oocyte maturation were obtained from the stages of GV, GVBD, metaphase I and metaphase II. Statistical analysis was performed by means of one-way Anova, followed by Tukey's HSD test. The level of significance was not at 0.05.

Ethical clearance

This study was given ethical clearance by the Research Ethics Committee, Faculty of

The number of oocytes was inversely proportional to the nicotine dose, i.e. larger doses of nicotine resulted in reduced number of oocytes.

The results of one-way Anova showed significant differences in the numbers of oocytes between all four groups ($p=0.000$). The control group had a significantly larger number of oocytes in comparison with the three treatment groups ($p=0.00$). The reduction in the number of oocytes in the group receiving nicotine at a dose of 21 mg/BW (I) was significantly smaller than that in the groups receiving nicotine at doses of 42 mg/kg BW (II) and 84 mg/kg BW (III) ($p=0.00$). The nicotine dose of 42 mg resulted in a smaller reduction in the number of oocytes than the dose of 84 mg, but the difference was statistically not significant ($p=0.74$).

The maturation of the oocytes was subsequently observed, with the results shown in Figure 2, depicting oocyte maturation stages in the control group and the groups exposed to nicotine at various doses. Exposure to nicotine at a dose of 21 mg/kgBW significantly reduced the numbers of oocytes in the GV stage as compared with controls ($p<0.05$). Exposure to doses of 42 and 84 mg/kgBW did not significantly alter the numbers of GV compared with controls ($p>0.05$). In the GVBD stage, exposure to nicotine at 21 mg/kgBW significantly reduced the number of oocytes in that stage as compared




Figure 1. Effect of nicotine exposure at various doses on number of mature oocytes in *Rattus norvegicus*




Figure 2. Effect of exposure to nicotine at various doses on oocyte maturation in *Rattus norvegicus*.

GV=germinal vesicle; GVBD=germinal vesicle breakdown

with controls ($p < 0.05$). Exposure to doses of 42 and 84 mg/kgBW did not significantly alter the numbers of oocytes in the GVBD stage as compared with controls ($p > 0.05$). In metaphase I, a significant reduction in the number of oocytes was found on exposure to nicotine doses of 42 and 84 mg/kgBW in comparison with controls ($p < 0.05$), which was not found with the dose of 21 mg/kgBW ($p > 0.05$). In metaphase II, a significant reduction in the number of oocytes was found on exposure to nicotine at all doses in comparison with controls ($p < 0.05$).

Tukey's HSD found significant differences between controls and the three nicotine dosage groups with respect to all observed maturation stages, with $p = 0.000$ for all comparisons.

For the GV stage, there were significant differences between controls and I, controls and II, and between I and II, I and III, II and III. For GVBD, the differing pairs of groups were controls and I, I and II, I and III. For metaphase I, the

Univ Med
32 No.2

differences were between controls on the one hand and I, II, and III on the other, and also between I and II, and between I and III. In metaphase II stage, the differing groups were controls versus I, II, and III, I and II, I and III.

DISCUSSION

In this study there were differences in the number of oocytes released from the ovaries between controls, treatment groups I (21 mg/kg BW), II (42 mg/kg BW) and III (84 mg/kg BW). Our study results are consistent with the study conducted by Mokhtar et al., where it was shown that nicotine adversely affects the number and quality of oocytes and the fertilization rate in animal models.⁽¹²⁾ A recent study using confocal microscopy on oocytes from mice exposed to cigarette smoke revealed that these oocytes had significantly thicker zona pellucida and shorter and wider meiotic spindles. Approximately 25% of these oocytes had errors in chromosomal congression or abnormally shaped spindles.⁽¹³⁾

In the present study, observations on the maturation stages of retrieved oocytes, comprising GV, GVBD, metaphase I and

Vol.

metaphase II, showed significant differences at all maturation stages between the control group and the three treatment groups. This study was an improvement over the study of Dwirahayu, who made similar observations with different doses. The study of Dwirahayu showed significant differences for metaphase II only.⁽⁸⁾ In contrast, for GV and GVBD there

were no significant differences, and for metaphase I no statistical tests could be performed because all results were zero.⁽⁶⁾ Another study using nontoxic doses of nicotine of 1.0, 2.5, 5.0 and 10.0 mmol/L, respectively, showed nicotine to have no adverse effects on GV breakdown.⁽¹⁴⁾

Free radical or reactive oxygen species (ROS) production by nicotine is the result of inhibition of anti-oxidant enzyme and subsequent lipid peroxidation. Oxidative stress from free radicals or ROS may damage the cell membrane and also induce DNA fragmentation.⁽¹⁵⁾ Oxidative stress also leads to chromosomal instability and programmed cell death, the latter being the main mechanism of oocyte death.

When fully developed or mature oocytes are released from the follicle for ovulation, the meiotic process is completed as shown by their being in metaphase II. If the meiotic process is not completed, it will stop at any given stage, either GV, GVBD or metaphase. If the developing oocytes have not reached their full size when released from the follicle, they cannot become mature, being in the GV and GVBD stages. Medium-sized oocytes may reach maturity although they have not yet completed the meiotic process and have stopped at metaphase I.⁽¹⁰⁾

The mechanism of oocyte maturation blockage as a result of exposure to nicotine is by systemic oxidative stress and oxidative stress in the follicular fluid. Intrafollicular oxidative stress may cause apoptosis of the follicular granulosa cells, thus impeding the process of folliculogenesis and reducing follicular diameter.⁽⁴⁾

The size of the follicle affects oocyte development because the capacity of oocytes to complete meiosis for full maturation depends on follicular size. If the follicle becomes smaller in size, the maturation process of the oocytes within the follicle will be blocked. The number of oocytes from small follicles are lower than that of mature oocytes from large follicles.^(16,17) Immature oocytes are characterized by the GV stage, GVBD and

metaphase I, whereas mature oocytes are characterized by metaphase II.^(16,17)

The results of the present study showed that each group differs significantly from the others, signifying that low doses of nicotine is capable of affecting oocyte maturational development. Increasing the dose severely affects the development resulting in fewer mature oocytes. Exposure to nicotine from cigarette smoke at any dose significantly affects oocyte development in metaphase I and II. If there are no oocytes in metaphase II, there are no mature oocytes. Ultimately, this results in female infertility because only mature oocytes can be fertilized by sperm. These findings demonstrate the extreme sensitivity of human oocytes to cigarette smoke, and underline the need for experimental animal data to clarify the causes of meiotic blockage.

One limitation of this study was the inability to perform nicotine exposure by inhalation in order to mimic smoking. The implication of the study is that the dose of 21 mg/kgBW, which was capable of reducing the number of oocytes in *Rattus norvegicus*, can be used to calculate the minimal dose in humans, i.e. 1176 mg. One cigarette contains 0.3-2 mg of nicotine, therefore 588-3920 cigarettes smoked actively or passively, are sufficient to impair fertility in women.

CONCLUSION

Nicotine, the major alkaloid in tobacco, is capable of blocking oocyte maturation in the Norway rat (*Rattus norvegicus*). Immature oocytes cannot be fertilized by sperm, invariably leading to infertility.

ACKNOWLEDGEMENT

We thank the staff at the Embryology Laboratory, Faculty of Veterinary Medicine, University of Airlangga, Surabaya and Bambang Setiawan, S. Ked., M. Biomed for enabling us to complete and publish this study.

REFERENCES

1. Rabinoff M, Caskey N, Rissling A, Park C. Pharmacological and chemical effects of



- cigarette additives. Am J Public Health 2007;97:1981-8.
2. Hukkanen J, Jacob P III, Benowitz NL. Metabolism and disposition kinetics of nicotine. Pharmacol Rev 2005;57:79-115.
3. Vasquez J. Human rights and health: persons exposed to second-hand tobacco smoke. Washington DC: Pan American Health Organization; 2008.
4. Bordel R, Laschke MW, Menger MD, Vollmar B. Nicotine does not affect vascularization but inhibits growth of freely transplanted ovarian follicles by inducing granulosa cell apoptosis. Hum Reprod 2006;21:610-6.
5. Talbot P, Riveles K. Smoking and reproduction: the oviduct as a target of cigarette smoke. Reprod Biol Endocrinol 2005;3:52. doi:10.1186/1477-78273-52.
6. Bordel R, Laschke MW, Menger MD, Vollmar B. Nicotine does not affect vascularization but inhibits growth of freely transplanted ovarian follicles by inducing granulosa cell apoptosis. Hum Reprod 2006;21:610-7.
7. Mailhes JB, Young D, Caldito G, London SN. Sensitivity of mouse oocytes to nicotine induced perturbations during oocyte meiotic maturation and aneuploidy in vivo and in vitro. Mol Hum Reprod 2000;6:232-7.
8. Dwirahayu Y. Pengaruh nikotin terhadap tingkat maturasi oosit pada tikus (*Rattus novergicus*) [thesis]. Surabaya: Universitas Airlangga; 2010.
9. Kakisina P. Pengaruh nikotin terhadap perkembangan embrio mencit (*Mus musculus*) [thesis]. Surabaya: Universitas Airlangga; 2003.
10. Kusumawati D. Bersahabat dengan hewan coba. Yogyakarta: Gadjah Mada University Press; 2004.
11. Shah H. How to calculate the sample size for animal studies.? Nat J Physiol Pharm Pharmacol 2011;1: 35-9.
12. Mokhtar N, Rajikin M, Zakaria Z. Role of tocotrienol-rich palm vitamin E on pregnancy and preimplantation embryos in nicotine-treated rats. Biomed Res 2009;19:181-4.
13. Jennings PC, Merriman JA, Beckett EL, Hansbro PM, Jones KT. Increased zona pellucida thickness and meiotic spindle disruption in oocytes from cigarette smoking mice. Hum Reprod 2011;26:878-84.
14. Zenzes MT, Bielecki R. Nicotine-induced disturbances of meiotic maturation in cultured mouse oocytes: alterations of spindle integrity and chromosome alignment. Tob Induced Dis 2004;2:151-61.
15. Asadi E, Jahanshahi M, Gholipour MJ. Effect of vitamin E on oocytes apoptosis in nicotine-treated mice. Iran J Basic Med Sci 2012;15:880-7.
16. Son WY, Lim JH. Fertilization, cleavage and blastocyst development according to the maturation timing of oocytes in invitro maturation cycles. Hum Reprod 2005;20:3204-7.
17. Mrazek M, Fulka Jr J. Failure of oocyte maturation: possible mechanisms for oocyte maturation arrest. Hum Reprod 2003;18:2249-52.